

Search Bioscience Technology:  

  
 Advanced Search

**Bioscience**  
 TECHNOLOGY.

[Log in](#) | [Register](#)

- [Directory](#)
  - [Publications](#)
  - [Sections](#)
  - [News](#)
  - [Products](#)
  - [Multimedia](#)
  - [Community](#)
- [Subscribe](#)

[Home](#) > [News](#)

- [RSS Feeds](#)
  - [Newsletters](#)
  - [Bookmarks](#)
- [\[-\] Text](#) [\[+\]](#)

## Forests absorb one-third of global fossil fuel emissions

Featured In: [Academia News](#)

By University of Leeds

Friday, July 15, 2011

[See today's top life science stories and headlines - Sign up now!](#)



The world's established forests remove 8.8bn tonnes of CO<sub>2</sub> per year from the atmosphere – equivalent to nearly a third of annual fossil fuel emissions – according to new research published in *Science*.

In addition, regrowth of trees on previously deforested lands in the tropics mopped up a further 6 billion tonnes of CO<sub>2</sub> annually between 1990 and 2007(1). However, deforestation across the tropics released a huge 10.8 billion tonnes of CO<sub>2</sub> annually during this period, off-setting much of the uptake of CO<sub>2</sub> by the world's forests. For comparison, global fossil fuel emissions average 28 billion tonnes of CO<sub>2</sub> annually.

The findings suggest that the world's tropical, temperate and boreal(2) forests play a much larger role in the global cycling of carbon than previously thought, and that protecting them is vital in limiting the severity of future climate change.

Dr Simon Lewis, a tropical ecologist from the University of Leeds and co-author of the study, said: "Humans are altering the world's forests in a number of ways, from their outright destruction to the much more subtle impacts on even the most remote forests caused by global changes to the environment.

"Our research shows these changes are having globally important impacts, which highlights the critical role forests play in the global cycling of carbon and therefore the speed and severity of future climate change.

"The practical importance of this new information is that if schemes to reduce deforestation are successful they would have significant positive global impacts, as would similar efforts promoting forest restoration."

The international team of researchers, led by Dr Yude Pan from the United States Forest Service, used on-the-ground measurements of trees worldwide and statistical models to provide an updated picture of atmospheric carbon uptake and loss by boreal, temperate and tropical forests spanning a total of 3.9 billion hectares.

Dr Pan said, "By breaking the large-scale carbon dynamics of forests into their many components, we were able to separately analyse the large magnitudes of carbon fluxes, both sinks and sources, and gain an idea of the potential capacity for carbon sequestration by the world's forests."

After adding together all the sources and sinks, the researchers found that there was a net climate benefit of 4 billion tonnes of carbon dioxide each year absorbed by forests, as uptake out-stripped releases.

The study also provided two further new insights.

In the tropics, separating the carbon losses from deforestation and carbon uptake from regrowing forests shows that the figure usually reported for carbon emissions associated with tropical 'land-use change' hides these two much larger fluxes. "By separating them out, we see that carbon emissions from tropical



Email Print

9 tweets

retweet

40

Share

0

Digg ↑

News

Supplier Update

[Appointment of Pro-Vice-Chancellor for Research and Innovation](#)  
 48 minutes ago

[Insurer Aetna to spend \\$202M on PayFlex Holdings](#)  
 1 hour ago

[OHSU scientists discover new role for vitamin C in the eye — and the brain](#)  
 2 hours ago

[Making Blood-Sucking Deadly for Mosquitoes](#)  
 2 hours ago

[Early-stage melanoma tumors contain clues to metastatic potential](#)  
 2 hours ago

[New technique boosts efficiency of blood cell production from human stem cells\(2\)](#)  
 2 hours ago

[Appointment for Professor David Hogg](#)  
 2 hours ago

BIOSCIENCE TECHNOLOGY Webcast Series

High-throughput long non-coding RNA gene expression profiling using qPCR

Sponsored by:

VIEW TODAY • REGISTER FREE

Most Popular

Related Content

[Penn Researchers Reprogram Brain Cells to Become Heart Cells](#)  
 Jul 11

deforestation are much larger than we thought, while uptake from forest re-growth in the tropics is also probably very large," said Dr Lewis.

The second insight is that the newly reported total carbon uptake by boreal, temperate and remaining tropical forests together is equal to the total global land carbon sink(3). "The large uptake of CO<sub>2</sub> by forests implies that the world's agricultural lands, grasslands, desert and tundra each play a more limited role as globally significant carbon dioxide sources or sinks at present. This new information can help pinpoint where actions to conserve carbon sinks are likely to have most impact," explains Dr Lewis.

However, Dr Lewis urged caution in the interpretation of the results: "We can't rely on forest management alone to halt the increase of CO<sub>2</sub> in the atmosphere and solve the problem of climate change. There is simply not enough land on Earth to store all the carbon released from ongoing fossil fuel emissions in trees.

"Forest management can help, but reducing fossil fuel emissions is essential. And, of course, forests won't keep removing CO<sub>2</sub> from the atmosphere forever. The world's remaining forests are providing a time-limited benefit to humanity."

Study co-author Oliver Phillips, Professor of Tropical Ecology at the University of Leeds, emphasised the need for more research. He said: "We know the tropics are the most dynamic area of the world when considering the exchange of carbon between the land and atmosphere.

"Trees grow fast in the tropics, and widespread deforestation is the norm, yet our collective research effort is smaller in the tropics than elsewhere. What we need is serious investment in monitoring the world's tropical forests to better understand their role in our rapidly changing global environment."

Professor Phillips co-ordinates the RAINFOR network of long-term tropical forest monitoring sites across South America (<http://www.rainfor.org/>). Dr Lewis co-ordinates a similar network across Africa (<http://www.afritron.org/>). Both were used to estimate the carbon sink in remaining tropical forests reported in the study.

For more information

The paper, '[A Large and Persistent Carbon Sink in the World's Forests, 1990-2007](#)', by: Pan Yude, Richard Birdsey, Jingyun Fang, Richard Houghton, Pekka Kauppi, Werner A. Kurz, Oliver L. Phillips, Anatoly Shvidenko, Simon L. Lewis, Josep G. Canadell, Philippe Ciais, Robert B. Jackson, Stephen Pacala, A David McGuire, Shilong Piao, Aapo Rautiainen, Stephen Sitch and, Daniel Hayes, is published in Science Express on 14 July 2011.

Contact

Hannah Isom  
Press Officer, University of Leeds  
T: +44 (0)113 343 5764  
E: [h.isom@leeds.ac.uk](mailto:h.isom@leeds.ac.uk)

[SOURCE](#)

Join the Discussion

Rate Article:  Average 0 out of 5

[register](#) or [log in](#) to comment on this article!

 0 COMMENTS

 ADD COMMENT

Text Only 2000 character limit

Page 1 of 1

## Research Exchange

Application  
Notes

White  
Papers

Posters

Journal  
Articles

Ask the  
Experts

Forests absorb one-third of global fossil fuel emissions

Jul 15

Cancer Gene Therapy from Camels

Jul 14

When Flu Strains 'Hook Up' Dangerous Progeny Can Result

Jul 14

Nutty Solution to Type 2 Diabetes Management


Jul 12

STAY INFORMED: SUBSCRIBE TO

**Bioscience  
TECHNOLOGY.**

Magazine and E-mail Newsletters

E-mail:



BIOSCIENCE TECHNOLOGY Webcast Series

**A Metabolomics Approach to the Study of Neurodegenerative Disease**

ON DEMAND NOW | Sponsored by **esa** & **DIONEX**

**REGISTER FREE**

Free Life Science Industry Subscriptions

Magazine

Newsletters



Sign up now ►

## MULTIMEDIA

Video:

Viewing SureFocus Slides  
Jun 11 2010